

Title	Efficacy and safety of treatment with biologicals (benralizumab, dupilumab, mepolizumab, omalizumab and reslizumab) for severe eosinophilic asthma
Authors	Agache, Ioana;Beltran, Jessica;Akdis, Cezmi;Akdis, Mubeccel;Canelo-Aybar, Carlos;Canonica, Walter;Casale, Thomas;Chivato, Tomas;Corren, Jonathan;Del Giacco, Stefano;Eiwegger, Thomas;Firin, Davide;Gern, James E.;Hamelmann, Eckard;Hanania, Nicola;Mäkelä, Mika;Martín, Irene Hernández;Nair, Parameswaran;O'Mahony, Liam;Papadopoulos, Nikolaos G.;Papi, Alberto;Park, Hae-Sim;Pérez de Llano, Luis;Posso, Margarita;Rocha, Claudio;Quirce, Santiago;Sastre, Joaquin;Shamji, Mohamed;Song, Yang;Steiner, Corinna;Schwarze, Jurgen;Alonso-Coello, Pablo;Palomares, Oscar;Jutel, Marek
Publication date	2020-02-08
Original Citation	Agache, I., Beltran, J., Akdis, C., Akdis, M., Canelo-Aybar, C., Canonica, W., Casale, T., Chivato, T., Corren, J., Del Giacco, S., Eiwegger, T., Firinu, D., Gern, J. E., Hamelmann, E., Hanania, N., Mäkelä, M., Martín, I. H., Nair, P., O'Mahony, L., Papadopoulos, N. G., Papi, A., Park, H.-S., Pérez de Llano, L., Posso, M., Rocha, C., Quirce, S., Sastre, J., Shamji, M., Song, Y., Steiner, C., Schwarze, J., Alonso-Coello, P., Palomares, O. and Jutel, M. (2020) 'Efficacy and safety of treatment with biologicals (benralizumab, dupilumab, mepolizumab, omalizumab and reslizumab) for severe eosinophilic asthma', <i>Allergy</i> . doi: 10.1111/all.14221
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://onlinelibrary.wiley.com/doi/abs/10.1111/all.14221 - 10.1111/all.14221
Rights	© 2020, EAACI and John Wiley and Sons A/S. Published by John Wiley and Sons Ltd. This is the peer reviewed version of the following article: Agache, I., Beltran, J., Akdis, C., Akdis, M., Canelo-Aybar, C., Canonica, W., Casale, T., Chivato, T., Corren, J., Del Giacco, S., Eiwegger, T., Firinu, D., Gern, J. E., Hamelmann, E., Hanania, N., Mäkelä, M., Martín, I. H., Nair, P., O'Mahony, L., Papadopoulos, N. G., Papi, A., Park, H.-S., Pérez de Llano, L., Posso, M., Rocha, C., Quirce, S., Sastre, J., Shamji, M., Song,

	Y., Steiner, C., Schwarze, J., Alonso-Coello, P., Palomares, O. and Jutel, M. (2020) 'Efficacy and safety of treatment with biologicals (benralizumab, dupilumab, mepolizumab, omalizumab and reslizumab) for severe eosinophilic asthma', Allergy, doi: 10.1111/all.14221, which has been published in final form at https://doi.org/10.1111/all.14221 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Download date	2023-05-05 02:36:36
Item downloaded from	http://hdl.handle.net/10468/9742



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

PROF. IOANA AGACHE (Orcid ID : 0000-0001-7994-364X)

MRS. MÜBECCEL AKDIS (Orcid ID : 0000-0002-9228-4594)

PROF. THOMAS B CASALE (Orcid ID : 0000-0002-3149-7377)

DR. THOMAS EIWEGGER (Orcid ID : 0000-0002-2914-7829)

DR. DAVIDE FIRINU (Orcid ID : 0000-0002-5768-391X)

PROF. ECKARD HAMELMANN (Orcid ID : 0000-0002-2996-8248)

DR. PARAMESWARAN NAIR (Orcid ID : 0000-0002-1041-9492)

PROF. LIAM O'MAHONY (Orcid ID : 0000-0003-4705-3583)

DR. NIKOLAOS G PAPADOPOULOS (Orcid ID : 0000-0002-4448-3468)

PROF. HAE-SIM PARK (Orcid ID : 0000-0003-2614-0303)

DR. LUIS PEREZ DE LLANO (Orcid ID : 0000-0003-2652-6847)

PROF. JOAQUIN SASTRE (Orcid ID : 0000-0003-4689-6837)

DR. MOHAMED H SHAMJI (Orcid ID : 0000-0003-3425-3463)

DR. OSCAR PALOMARES (Orcid ID : 0000-0003-4516-0369)

Article type : Guidelines

Corresponding author email id: ibrumaru@unitbv.ro

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/all.14221](https://doi.org/10.1111/all.14221)

This article is protected by copyright. All rights reserved

Efficacy and safety of treatment with biologicals (benralizumab, dupilumab, mepolizumab, omalizumab and reslizumab) for severe eosinophilic asthma

A systematic review for the EAACI Guidelines - recommendations on the use of biologicals in severe asthma

Authors

Ioana Agache^{*1}, Jessica Beltran^{*2}, Cezmi Akdis³, Mubeccel Akdis³, Carlos Canelo-Aybar^{2,4}, Walter Canonica⁵, Thomas Casale⁶, Tomas Chivato⁷, Jonathan Corren⁸, Stefano Del Giacco⁹, Thomas Eiwegger^{10,11,12}, Davide Firinu⁹, James E. Gern¹³, Eckard Hamelmann¹⁴, Nicola Hanania¹⁵, Mika Mäkelä¹⁶, Irene Hernández Martín¹⁷, Parameswaran Nair¹⁸, Liam O'Mahony¹⁹, Nikolaos G. Papadopoulos^{20,21}, Alberto Papi²², Hae-Sim Park²³, Luis Pérez de Llano²⁴, Margarita Posso^{2,25}, Claudio Rocha², Santiago Quirce²⁶, Joaquin Sastre²⁷, Mohamed Shamji^{28,29}, Yang Song², Corinna Steiner², Jurgen Schwarze³⁰, Pablo Alonso-Coello^{**2, 4}, Oscar Palomares^{**31}, Marek Jutel^{**32,33}

*Joint first authorship

** Joint last authorship

Corresponding author: Ioana Agache; 2A, Pictor Ion Andreescu, Brasov, Romania, 500051

Affiliations

¹Transylvania University, Faculty of Medicine, Brasov, Romania

²Iberoamerican Cochrane Centre - Department of Clinical Epidemiology and Public Health, Biomedical Research Institute Sant Pau (IIB Sant Pau), Barcelona, Spain

³ Swiss Institute of Allergy and Asthma Research (SIAF), University of Zurich, Davos, Switzerland

⁴CIBER de Epidemiología y Salud Pública (CIBERESP), Spain

⁵ Personalized Medicine, Asthma and Allergy, Humanitas Clinical and Research Center, IRCCS, Rozzano, Italy.

⁶ Division of Allergy and Immunology, University of South Florida Morsani College of Medicine, Tampa, FL, USA

⁷ School of Medicine University CEU San Pablo, Madrid, Spain

⁸ David Geffen School of Medicine at UCLA, Los Angeles, USA

⁹ Department of Medical Sciences and Public Health, University of Cagliari, Italy

- ¹⁰ Translational Medicine Program, Research Institute, Hospital for Sick Children, Toronto, ON, Canada
- ¹¹ Department of Immunology, University of Toronto, Toronto, ON, Canada
- ¹² Division of Immunology and Allergy, Food Allergy and Anaphylaxis Program, The Hospital for Sick Children, Departments of Paediatrics and Immunology, University of Toronto, Toronto, Canada
- ¹³ Department of Pediatrics; University of Wisconsin School of Medicine and Public Health
- ¹⁴ Klinik für Kinder- und Jugendmedizin Kinderzentrum Bethel
- ¹⁵ Section of Pulmonary, Critical Care and Sleep Medicine, Baylor College of Medicine, Houston, Texas, USA
- ¹⁶ Skin and Allergy Hospital, Helsinki University Hospital and University of Helsinki, Finland
- ¹⁷ Department of Allergy, Hospital Universitario La Paz, Madrid, Spain
- ¹⁸ Division of Respiriology, Department of Medicine, McMaster University, Hamilton, ON, Canada; Firestone Institute for Respiratory Health, St Joseph's Healthcare, Hamilton, ON, Canada
- ¹⁹ Departments of Medicine and Microbiology, APC Microbiome Ireland, University College Cork, Cork, Ireland
- ²⁰ Division of Infection, Immunity & Respiratory Medicine, University of Manchester, Manchester, UK
- ²¹ Allergy Dpt, 2nd Pediatric Clinic, National Kapodistrian University of Athens, Athens, Greece
- ²² Research Center on Asthma and COPD, Department of Medical Sciences, University of Ferrara, Ferrara, Italy
- ²³ Department of Allergy and Clinical Immunology, Ajou University, South Korea
- ²⁴ Department of Respiratory Medicine, Hospital Lucus Augusti, Lugo, Spain
- ²⁵ Department of Epidemiology and Evaluation, IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain.
- ²⁶ Department of Allergy, La Paz University Hospital, IdiPAZ, CIBER of Respiratory Diseases (CIBERES), Universidad Autónoma de Madrid, Spain
- ²⁷ Universidad Autónoma de Madrid Facultad de Medicina
- ²⁸ Immunomodulation and Tolerance Group, Allergy and Clinical Immunology, Inflammation, Repair, Development, National Heart and Lung Institute, UK
- ²⁹ Imperial College NIHR Biomedical Research Centre, Asthma UK Centre in Allergic Mechanisms of Asthma, London, United Kingdom
- ³⁰ Centre for Inflammation Research, Child Life and Health, The University of Edinburgh, Edinburgh, United Kingdom
- ³¹ Department of Biochemistry and Molecular Biology, Chemistry School, Complutense University of Madrid, Spain
- ³² University of Wroclaw, Department of Clinical Immunology, Wroclaw Poland
- ³³ "ALL-MED" Medical Research Institute, Wroclaw, Poland

ABSTRACT

Five biologicals have been approved for severe eosinophilic asthma, a well-recognised phenotype. Systematic reviews (SR) evaluated the efficacy and safety of benralizumab, dupilumab, mepolizumab, omalizumab and reslizumab (alphabetical order) compared to standard of care for severe eosinophilic asthma.

Pubmed, EMBASE and Cochrane Library were searched to identify RCTs and health economic evaluations, published in English. Critical and important asthma-related outcomes were evaluated for each of the biologicals. The risk of bias and the certainty of the evidence were assessed using GRADE.

19 RCTs (3 RCTs for benralizumab, 3 RCTs for dupilumab, 3 RCTs for mepolizumab, 4 RCTs for omalizumab, and 5 RCTs for reslizumab), including subjects 12 to 75 years old (except for omalizumab including also subjects 6-11 years old), ranging from 12 to 56 weeks were evaluated. All biologicals reduce exacerbation rates with high certainty of evidence: benralizumab rate ratio (RR) 0.53 (95% CI 0.39 to 0.72), dupilumab RR 0.43 (95% CI 0.32 to 0.59), mepolizumab RR 0.49 (95% CI 0.38 to 0.66), omalizumab RR 0.56 (95% CI 0.40 to 0.77), and reslizumab RR 0.46 (95% CI 0.37 to 0.58). Benralizumab, dupilumab and mepolizumab reduce the daily dose of oral corticosteroids (OCS) with high certainty of evidence. All evaluated biologicals probably improve asthma control, QoL and FEV₁, without reaching the minimal important difference (moderate certainty). Benralizumab, mepolizumab and reslizumab slightly increase drug-related adverse events (AE) and drug-related serious AE (low to very low certainty of evidence). The incremental cost-effectiveness ratio per quality adjusted life-years value is above the willingness to pay threshold for all biologicals (moderate certainty). Potential savings are driven by decrease in hospitalisations, emergency and primary care visits.

There is high certainty that all approved biologicals reduce the rate of severe asthma exacerbations and for benralizumab, dupilumab and mepolizumab for reducing OCS. There is moderate certainty for improving asthma control, QoL, FEV₁ and cost-effectiveness. More data on long term safety are needed together with more efficacy data in the paediatric population.

Abbreviations

ACQ = asthma control questionnaire

AQLQ = asthma related quality of life questionnaire

AE = adverse events

CHEC = Consensus health economic criteria

CI = confidence interval

EAACI = European Academy of Allergy and Clinical Immunology

EMA = European Medicine Agency

EURONHEED = European Network of Health Economic Evaluation Databases

FDA = Food and Drug administration

FeNO = fractional exhaled nitric oxide

FEV₁ = forced expiratory volume in one second

GDG = Guideline Development Group

GINA = Global Initiative for Asthma

GRADE = Grading of Recommendations Assessment, Development and Evaluation

ICS = inhaled corticosteroids

ICER = Incremental cost-effectiveness ratio

Ig = immunoglobulin

IL = interleukin

IRR = incidence rate ratios

IV= intravenous

MD = mean difference

MID = minimal important difference

OCS = oral corticosteroids

QALY = Quality adjusted life-years

QoL = quality of life

RCT = randomised controlled trial

ROB = risk of bias

SGRQ = St George's Respiratory Questionnaire

SOC= standard of care

RR= rate ratio

SC= subcutaneous

SR = systematic review

TASS = Total Asthma Symptoms Scores

T2 = type 2

Accepted Article

Key words

severe asthma; cost-effectiveness; eosinophilic asthma; exacerbations; benralizumab; dupilumab; mepolizumab; omalizumab; reslizumab

Introduction

There are still a significant proportion of patients with severe asthma having uncontrolled or partially controlled asthma despite correct management. These patients represent a special challenge related to extensive diagnostic evaluation and high consumption of health-care resources (1,2,3,4). Severe asthma is defined as inadequate control of asthma under high-intensity treatment with inhaled corticosteroids (ICS) and additional controllers (including oral corticosteroid (OCS)) for at least six months per year, or by loss of asthma control on the attempt to reduce the high-intensity treatment (5). Before any further treatments are evaluated, differential diagnoses of asthma should be ruled out, comorbidities should be treated, persistent triggers should be eliminated, and patient adherence should be optimised (2,5,6). Considering the availability of specific targeted therapies for type 2 (T2) asthma, the management approach to severe asthma currently includes a phenotyping step for the identification of allergic, eosinophilic and non-T2 phenotypes (7,8).

Eosinophils are prominent pathogenic cells involved in asthma. Increased blood or sputum eosinophils were related to frequent asthma exacerbations and disease severity and are used to guide treatment decisions (9,10,11). Most human diseases accompanied by elevated blood eosinophils are associated with increased interleukin (IL)-5 production. Although IL-5 plays a central role in eosinophil biology, it is neither necessary nor sufficient for fully inducing an eosinophil mediated disease. In humans, IL-5 is often co-expressed with other T2 cytokines including IL-4 and IL-13 and associated in atopic individuals with increased immunoglobulin (Ig)-E production (12,13,14,15,16). This over-lap between the key pathogenetic pathways (IL-5, IL-4/IL-13 and IgE driven eosinophilic inflammation) toughen the choice of a biological for the eosinophilic asthma phenotype. Therefore, future novel approaches to better identify predominant pathways at the molecular level (endotypes) are demanded.

In the last decades, new add-on therapies for severe asthma have been developed, and may be applied depending on asthma phenotype and endotype (7, 8, 13). Anti-IL-5 monoclonal antibodies (mepolizumab and reslizumab) are approved for severe asthma and peripheral eosinophilia (17,18,19,20). Benralizumab, a monoclonal antibody that binds to the α subunit of IL-5 receptor (IL-5R α) was also recently approved for eosinophilic asthma (21,22). Dupilumab, a monoclonal antibody directed against the α subunit of the IL-4 receptor (IL-4R α) acting as a dual inhibitor of both IL-4- and IL-13-mediated signalling pathways was approved for T2 asthma (23,24). Omalizumab, a humanised monoclonal anti-IgE antibody, was the first biological approved for IgE-mediated persistent allergic asthma (25,26).

The European Academy of Allergy and Clinical Immunology (EAACI) is developing clinical practice guidelines for the use of biologicals in patients with severe asthma. Three interlinked

systematic reviews (SR) were performed to inform the formulation of key clinical recommendations. The current SR is focusing on eosinophilic asthma assessing the current evidence regarding efficacy, safety and economic impact of the biologicals with current regulatory approval for patients with uncontrolled severe asthma (i.e. benralizumab, dupilumab, omalizumab, mepolizumab, and reslizumab, in alphabetical order).

METHODS

Guideline Development Group

The EAACI Asthma Voting Panel and Guidelines Steering Committee includes clinicians and researchers with different backgrounds (the complete list of experts is available from the EAACI website) whom voluntarily participate in the development of EAACI clinical practice guidelines for the use of biologicals in severe asthma. They are referred to as the Guideline Development Group (GDG).

Structured question and outcome prioritisation

The GDG framed the clinical question as: "Is the treatment with biologicals (i.e., benralizumab, dupilumab, mepolizumab, omalizumab and reslizumab) efficacious and safe for patients with uncontrolled severe eosinophilic asthma?" For the purpose of this SR the population of interest was defined as subjects with any of the following: a sputum eosinophil count of $\geq 1\%$ or an asthma-related peripheral blood eosinophil count of ≥ 150 cells/ μL , or a fractional exhaled nitric oxide (FeNO) of ≥ 20 ppb (27). The outcomes were prioritised by the GDG using a 1 to 9 scale (7 to 9 critical; 4 to 6 important; 1 to 3 of limited importance), as suggested by the GRADE approach. The critical outcomes were exacerbations, asthma control, QoL, and safety and the important outcomes included lung function (forced expiratory volume in one second – FEV₁), OCS and ICS use and rescue medication use (table S1). Safety was assessed evaluating the rate of drug-related adverse events (AE) or drug-related serious AEs.

The GDG also framed a cost-effectiveness question to assess the economic impact of these biologicals versus standard of care for patients with uncontrolled severe eosinophilic asthma. The outcomes of interest were costs and resource use, as well as the incremental cost-effectiveness ratios (ICERs) per both quality adjusted life-years (QALY) and per asthma-specific outcomes.

Data sources and searches

MEDLINE (via PubMed, January 2019), EMBASE (via Ovid, January 2019) and CENTRAL (via The Cochrane Library, January 2019) databases were searched using pre-defined algorithms for individual studies for evidence of effects and economic evaluations, including systematic reviews as source of individual studies. Search terms were adapted to each database, validated filters were used to retrieve appropriate designs. Members of the GDG were requested to provide additional studies for evaluation (Table S2).

Study selection

Randomised controlled trials (RCTs) of patients with uncontrolled severe eosinophilic asthma that compared benralizumab, dupilumab, mepolizumab, omalizumab or reslizumab versus standard of care were included in the SR. Separate searches for each of the five biologicals evaluated were performed. The SR excluded studies in which the dose or the route of the biological was not

approved by the European Medical Agency (EMA) or by U.S. Food and Drug administration (FDA). Abstracts or conference communications not published as full articles in peer reviewed journals were also excluded. Only studies published in English were considered. Two reviewers independently assessed the references based on title and abstract. Then, two reviewers independently assessed the eligibility of the studies according to inclusion criteria. Discrepancies were solved either by consensus or with the help of a third reviewer. All citations retrieved were imported into the bibliographic reference software (EndNote X5; Thomson Reuters) to discard duplicates and record screening decisions.

Data extraction and risk of bias assessment

Details of the study design, patient population, setting, follow-up, and results were extracted by one reviewer, and confirmed by a second reviewer. If needed, requested additional data from the authors of the included studies was requested. The risk of bias (ROB) was assessed using the Cochrane Risk of Bias Assessment tool (28). The ROB was judged as low, high or unclear risk for each domain: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, and selective reporting.

For the health economics analysis, two reviewers extracted the main characteristics of included studies (e.g. type of economic evaluation, perspective, time horizon, discounting, sources of information, model type), relevant outcomes and costs (e.g. ICERs, sensitivity analyses results), sources of funding, and conflict of interest. Two reviewers assessed the methodological limitations using the consensus on health economics criteria (CHEC) checklist (29). Transferability to the European context was assessed using the European Network of Health Economic Evaluation Databases (EURONHEED) checklist (30,31).

Data synthesis and analysis

The main results of the SR are described narratively and tabulated as summary of findings. For dichotomous variables data were pooled as incidence rate ratios (IRR) and risk ratios (RR). For continuous data mean differences, with 95% confidence intervals were used. Change in the observed asthma related outcomes were assessed between biologicals and placebo from baseline to the end of the treatment. A random-effects model was used to pool data (Review Manager V5.3). Where multiple arms were compared to a common placebo arm standard errors were adjusted to avoid the unit of analysis error (32).

Statistical heterogeneity between studies was assessed using the Cochrane chi-square test, and the magnitude of heterogeneity with the I^2 statistic. To account for clinical heterogeneity, the analyses were stratified whenever possible by different doses of the biological, age, relevant biomarkers (blood eosinophils, FeNO) and ROB. Additionally, post hoc subgroup analysis by

baseline use of OCS was performed. To estimate the absolute effects for each comparison the median estimate reported in the control arms was used as baseline risk. For the economic evidence, results are summarised narratively and tabulated, including the ICERs and the degree of uncertainty.

Certainty of evidence

The certainty (quality) of the evidence of efficacy, safety and economic impact for each outcome was rated as high, moderate, low or very low considering the standard GRADE domains: ROB, imprecision, inconsistency, indirectness and publication bias (33,34). For the evaluation of imprecision for each asthma related outcome evaluated, where available, the minimal important difference (MID) thresholds were considered (35, 36,37, 38). For FEV₁, a MID of 0.20 L was considered as recommended by consensus by the GDG.

RESULTS

Search results

The eligibility process is summarized in a PRISMA flowchart (Figure 1). A total of 3441 unique citations were retrieved from data base searches, and 145 were appraised as full text. 135 publications were excluded due to differences in population, outcomes of interest, design or regulatory unapproved dose and/or route (Table S3). 28 publications from 19 RCTs were evaluate. These included 3 RCTs for benralizumab (39,40,41); 3 for dupilumab (42,43,44); 3 for mepolizumab (45,46,47); 4 for omalizumab (48,49,50,51); and 5 for reslizumab (52,53,54,55).

Characteristic of included studies

The characteristics of studies evaluated for evidence of efficacy and safety are detailed in Table 1. All studies randomised to either an intervention arm or a standard of care/placebo arm. All were RCTs, conducted during the previous eight years (2011-2019). The follow up under study medication ranged from 12 to 56 weeks. All studies included subjects aged 12 to 75 years old and studies of omalizumab also included children from 6 years old. The characteristics of the health economics studies included are available in Table 2.

Evidence of efficacy and safety

Tables 3 to 7 present the summary of the results and certainty of evidence for asthma related outcomes evaluated. The metanalysis plots can be found in the supplementary file.

Severe asthma exacerbation rate

The annualized exacerbation rates were reported in 3 benralizumab trials (39,40,41); 3 dupilumab trials (42,43,44), 3 mepolizumab trials (45,46,47), 3 omalizumab trials (48,50,51) and 5

reslizumab trials (52,53,54,55). All biologicals reduced asthma exacerbations rate compared to standard of care with high certainty of evidence: benralizumab RR 0.53; 95% CI 0.39 to 0.72; dupilumab RR 0.44; 95% CI 0.32 to 0.59; mepolizumab RR 0.49 95% CI 0.38 to 0.66; omalizumab RR 0.56; 95% CI 0.40 to 0.77; and reslizumab RR 0.46; 95% CI 0.37 to 0.58.

A separate analysis of the studies (41,44,46) designed to assess as primary outcome the OCS sparing effect of the biological compared to standard of care was performed. All three studies significantly reduced the rate of exacerbations in the OCS sparing protocol with high certainty of evidence: benralizumab RR 0.30; 95% CI 0.17 to 0.53; dupilumab RR 0.42; 95%CI 0.25 to 0.69; and mepolizumab RR 0.68; 95%CI 0.47 to 0.98.

Asthma control

The change in asthma control following biologicals addition was evaluated using Asthma Control Questionnaires (ACQ) scores and the Total Asthma Symptoms Scores (TASS). Dupilumab, omalizumab and mepolizumab probably improve asthma control with moderate certainty of evidence: dupilumab (ACQ-5) MD -0.48; 95% -0.88 to -0.09 (42,43,44); omalizumab (TASS) MD -0.16; 95% -0.51 to 0.19 (48,49,50,51) and mepolizumab (ACQ-5) MD -0.43; 95% CI -0.56 to -0.31 (45,46,47). Nevertheless, none of the biologicals showed an improvement above the MID threshold of 0.5.

Quality of life

QoL was reported in 3 benralizumab trials (39,40,41); 2 dupilumab trials (42,43); 3 mepolizumab trials (45,46,47); 1 omalizumab trial (48) and 3 reslizumab trials (53, 54, 55). Changes in QoL were evaluated using the Asthma Quality of Life Questionnaire (AQLQ) for all biologicals, except for mepolizumab that used the St. George's Respiratory Questionnaire (SGRQ) score. All The addition of all biologicals improved QoL with moderate to high certainty, although below the MID: benralizumab MD +0.23 (95% CI 0.11 to 0.36); dupilumab MD +0.42 (95% CI +0.25 to +0.59); mepolizumab (SGRQ) MD -7.14 (95% CI -9.07 to -5.21); omalizumab MD +0.13 (95% CI +0.11 to +0.37); and reslizumab MD +0.17 (95% CI +0.08 to +0.25).

Safety

Drug-related AE were assessed in 2 trials for benralizumab (40,41), 1 trial for dupilumab (42), 3 mepolizumab trials (45,46,47), 1 trial for omalizumab (48) and 3 trials for reslizumab (52,53,55). For mepolizumab there is an increased likelihood of drug-related AE (RR 1.35; 95% CI 1.01 to 1.80; high certainty of evidence. Benralizumab and reslizumab probably increases drug-related AE (moderate certainty of evidence): benralizumab RR 1.41, 95% CI 0.87 to 2.27; reslizumab RR

1.18, 95% CI 0.89 to 1.56. For dupilumab and omalizumab the RR is rather small: dupilumab RR of 1.00, 95% CI 0.88 to 1.13; omalizumab RR 1.01, 95% CI 0.91 to 1.1.

There is low to very low certainty of evidence that drug-related serious AE may increase with the use of dupilumab RR 1.46 (95% 0.60 to 3.54) and reslizumab RR 4.71 (95% 0.54 to 41.31). For benralizumab and mepolizumab results are inconclusive: benralizumab RR 0.56 (95% CI 0.22 to 1.44); mepolizumab RR 0.98 (95% CI 0.06 to 15.63). Data were not fully reported in all trials thus the certainty of evidence was downgraded due to the low number of events.

Reduction of oral corticosteroids use

Benralizumab, dupilumab and mepolizumab showed with high certainty of evidence a reduction in daily OCS: benralizumab > 50% (RR 1.76, 95%CI 1.26 to 2.47; dupilumab 29.4% (95% CI 43.2 lower to 15.57 lower); and mepolizumab > 50% (RR 1.61; 95%CI 1.07-2.41) (41,44,46). Mepolizumab showed a reduction of OCS to 5mg/day or less (crude RR 1.71; 95%CI 1.11 to 2.55, p=0.01) and a reduction of 100% in daily OCS (crude RR 1.91; 95% CI 0.69 to 5.30, p=0.2) compared to placebo.

Reduction of rescue medication use

This end-point was assessed only for mepolizumab and showed no clinically significant reduction in the daily use of rescue medication after 24 weeks (MD - 0.1 puffs /day; CI 95% -0.35 to 0.15) (45).

Lung function - FEV₁

The change from baseline of FEV₁ were assessed for benralizumab (39,40,41), mepolizumab (45,46,47), omalizumab (48) and reslizumab (52,53,54,55). Compared to standard of care, there was an increase of FEV₁, but below the MID agreed by the GDG (moderate certainty of evidence): benralizumab MD +140mL (95% CI +90 to +190); mepolizumab MD +110.9 mL (95% CI +58.91 to +162.89), reslizumab MD +141.82 mL (95% CI +89.23 to +194.41); and omalizumab mean percentage change + 3.7% (95% CI 2.1% to 9.5%). There is low certainty of evidence that for patients with baseline eosinophils ≥300 cells/μl dupilumab may increase FEV₁ compared to standard of care [MD +180mL (95% CI 110 to 250)].

Evidence of resource use and cost-effectiveness

After screening 1884 hits and reviewing 36 full text articles, five economic evaluations were included (Figure 2). Two studies evaluated benralizumab (56,57), 1 dupilumab (57), 1 mepolizumab (58) and 2 reslizumab (59,60) (Table 2). Most of excluded studies evaluated

allergic patients (Table S4). Overall, the resources needed for adding the biological to standard asthma therapy are mainly the cost of the drug and its administration. The potential savings are related to decreased rate of hospitalisation, emergency department care, and primary care visits (Table S5). The evaluation of the cost-effectiveness of benralizumab shows important variation in ICER from 39,135 £/QALY (low certainty of the evidence) to 412,000\$/QALY (moderate certainty of the evidence). The key driver for this difference is unclear since there is missing information in the Tikhonova et al report (Table S6). For dupilumab the reported ICER is 401,000 \$/QALY in patients with ≥ 300 eosinophils/ μ L (moderate certainty of the evidence). The results are uncertain for the utility estimates for the non-exacerbation health state for both the biological and standard of care asthma therapy, for the annual exacerbation rates for standard therapy and for the costs of chronic OCS use (Table S6). For mepolizumab the cost utility Markov model with low risk of bias (high quality study) reported an ICER/QALY value of 385,546 \$ (low certainty of the evidence). There is also variation across patients. The most favourable ICER towards mepolizumab was 160,000 \$ in the responder group of patients (Table S6). The reslizumab base-case analyses demonstrate important variation across studies in terms of the cost-effectiveness results. The ICER/QALY varied from 23,081 \$/ QALY (low certainty of the evidence) to 697,403 \$ / QALY (moderate certainty of the evidence). The key driver for this difference was the cost of reslizumab. The study not funded by industry (60) reported a higher cost of reslizumab (approximately 4,000 \$ per month) compared to the study funded by industry (991 \$ per month) (59). It has to be noted that there is a lack of proper studies to drive firm conclusions due to the heterogeneity of published studies. Furthermore, modelling cost-effectiveness studies considering direct and indirect costs applicable to each context are needed (Table S6).

DISCUSSION

Main findings

Our systematic review of efficacy shows high certainty for reducing the rate of severe asthma exacerbations for all the biologicals evaluated (benralizumab, dupilumab, mepolizumab, omalizumab, reslizumab) as add-on treatment for patients with severe uncontrolled eosinophilic asthma. The certainty is moderate for improving asthma control, QoL and lung function (FEV₁) improvement, not reaching the MID. Only benralizumab, dupilumab and mepolizumab provided data about the use of OCS, showing a reduction in the daily dose of OCS compared to standard of care (high certainty of evidence).

The main reasons to downgrade the certainty of evidence were ROB, imprecision, and indirectness. Overall the included studies for the evidence of efficacy were of low ROB. Of note, all included studies were funded by industry and all showed positive results, which might raise concerns of on a potential sponsorship bias. Although short term safety data are reassuring, the rate of drug related AE was not comprehensively reported, thus there is low to very low certainty for drug-related serious AE. With the exception of omalizumab there is scarce data on the efficacy of the evaluated biologicals for eosinophilic asthma in the paediatric population. With the exception of omalizumab and mepolizumab no data are available for long term safety, both in children and in adults.

The resources needed for adding the biological treatment to asthma standard therapy are mainly driven by the cost of the drug and its administration. With the approval of autoinjectors for most of these agents the costs for administration should decline. The potential savings are explained by the decreased rate of hospitalisation, emergency department care, and primary care visits. However, we cannot accurately assess the reductions in indirect costs such as improvements leading to improved work and school productivity and decreased absenteeism. The current SR of cost-effectiveness showed, for all the biologicals, an ICER per QALY value significantly above the willingness to pay threshold in most European countries (30,000 €/QALY). The certainty of the economic evidence is moderate derived from studies with low ROB but with important imprecision (large variations in the ICER values) and indirectness (differences in healthcare systems and year of conduction of the study), impeding both the transferability and generalization of the results. Additional publication bias might be envisaged.

Current results in the context of previous results

Aligned with our results, previous SR of benralizumab, dupilumab, mepolizumab, omalizumab and reslizumab in severe asthma reported a reduction to approximately half of exacerbations, and an improvement in health related QoL scores, asthma control and FEV₁ (61, 62, 63, 64, 65, 66, 67, 68). However, there are some important differences that need to be mentioned. Previous mepolizumab SR allowed the inclusion of non-approved doses (64,65), while, the current SR included only licensed doses and routes of administration making the results more applicable to the daily practice. Another important difference is that the certainty of the evidence was assessed using the GRADE approach. In contrast with the previous reviews assessing only the ROB of the trials, the current SR evaluated the heterogeneity, imprecision and indirectness of the included evidence. As an example, using the MID to assess for the imprecision, one can appreciate the clinical relevance of the change (i.e. asthma control, QoL and FEV₁): although the change was statistically significant it did not reach the MID. Last but not least, the current review assessed the

effect of all the biologicals across a comprehensive compilation of asthma related outcomes giving the clinicians a better perspective of the drug profile.

In contrast with our results, a SR of 8 benralizumab trials showed a lower risk of both AE (RR 0.94; 95% CI 0.90–0.98) and serious AE (RR 0.82; 95% CI 0.68–0.98) (61). The possible explanation for this difference is that we limited our assessment to drug-related AE and excluded asthma worsening events, assessed solely as efficacy measures.

Another review assessing the cost-effectiveness of mepolizumab for eosinophilic asthma showed that mepolizumab is cost-effective only when targeting specific subgroups with very severe asthma or by considerable price discounts. The key drivers of cost-effectiveness included day-to-day health-related QoL, asthma-related mortality, acquisition price of the biological therapy, and time horizon. These results are similar to our evaluation for mepolizumab (69).

Strengths and limitations

The current systematic review has several strengths. An exhaustive evaluation of both desirable and undesirable effects of the use of biologicals, as well as their economic impact was performed. The compilation of critical and important asthma related outcomes provides a more comprehensive perspective of the drug profile. The current SR used rigorous methods including the GRADE approach to rate the certainty of the evidence, with transparent judgements about the quality of evidence. We included the most updated results available from the included RCTs and only considered licensed doses and routes of administration. We provided friendly tabulated summaries of findings using optimal presentation format for patients, clinicians and policy makers supporting all the stakeholders in their endeavour to formulate recommendations for the use of biologicals in severe uncontrolled eosinophilic asthma.

There are several limitations. The basal exacerbation rate was used to estimate the absolute benefit for each drug/analysis. However, we did not perform a subgroup or sensitivity analysis based on that variable (basal exacerbation rate), as it was not pre-defined or requested in the protocol or during the systematic review. The systematic review included only English language articles, however the risk of selection bias is probably small because we screened previous systematic reviews and the GDG included several international experts in the field, thus the possibility of missing results from non-English articles is unlikely. We did not include observational studies that could have been informative for some of the outcomes with low or very low-quality evidence from RCTs (e.g. serious AE). We did not conduct a *de novo* economic analysis for the cost-effectiveness outcomes. However, we followed the global perspective on the use of biological treatment in different health systems, which may be useful for the decision of using the biologicals in different countries. Finally, this review is limited to patients with

eosinophilic asthma which restricted the number and scope of studies analysed, especially for omalizumab and dupilumab.

Implications for practice and research

All evaluated biologicals (benralizumab, dupilumab, mepolizumab, omalizumab, reslizumab) showed a significant improvement in critical and important related asthma outcomes such as exacerbation rate and for benralizumab, dupilumab and mepolizumab also an OCS dose reduction. However, the systematic review confirms the dissociated effect as these drugs have a modest effect on asthma control, quality of life and lung function. The health-economics and ICER analyses demonstrate above the willingness to pay threshold. Given the high cost of these drugs their use will probably be limited to very specific circumstances such as patients with severe uncontrolled asthma where the desired outcome is to decrease the exacerbation rate or the OCS use. In this context, panels are more likely to formulate conditional recommendations as opposed to strong.

Although short term safety data is reassuring, more accurate reporting from clinical trials is warranted in combination with long-term safety and cost-effectiveness evaluations, including real-world studies, registries and big data analysis. There is limited data available to support the efficacy and safety in the paediatric population (with the exception of omalizumab), highlighting the urgent unmet need for rigorous trials with biologicals in severe eosinophilic asthma in this population. Finally, the better understanding of predominant and personalized asthma endotypes and well-controlled head-to-head, industry-independent comparisons would provide useful data to better inform clinicians about choosing the right biologic for the right patient.

CONFLICT OF INTEREST:

IA serves as associate editor of Allergy. **JB** reports funding from EAACI. **CA** reports grants from Allergopharma, Idorsia, Swiss National Science Foundation, Christine Kühne-Center for Allergy Research and Education, European Commission Horizon 2020 Framework Programme, Cure, Novartis Research Institutes, Astra Zeneca, Scibase, and is on the Sanofi/Regeneron advisory board. **CC-A** reports funding from EAACI. **T Casale** reports grants and/or personal fees from Genentech, Novartis, Sanofi Regeneron, and GSK. **JC** declares grants or personal fees from AZ, Genentech/Roche, Novartis, Optinose, Sanofi, Stallergenes, and TEVA. **SG** reports personal fees from AstraZeneca, GSK and Novartis. **TE** has received grants or other from DBV, Innovation Fund Denmark, Regeneron, the Allergy and Anaphylaxis Program SickKids; serves as associate editor

for Allergy and in the local advisory board of ALK. **JG** reports personal fees from Regeneron, Ena Therapeutics, MedImmune/AstraZeneca, and stock options from Meissa Vaccines Inc. **NH** reports funding, honoraria or personal fees from GSK, AstraZeneca, Boehringer Ingelheim, Novartis and Sanofi Genzyme, Regeneron, Genentech, Sunovion, and Mylan. **PN** reports grants and/or personal fees from AZ, Novartis, Teva, Sanofi, Roche, Novartis, Merck, and Equillium. **LOM** reports grants from GSK and personal fees from AHL. **NP** reports grants from Gerolymatos and personal fees from Hal Allergy B.V., Novartis Pharma AG, Menarini, Hal Allergy B.V., and Mylan. **AP** has received grants, personal fees, non-financial support or other from GlaxoSmithKline, AstraZeneca, Boehringer Ingelheim, Chiesi Farmaceutici, TEVA, Mundipharma, Zambon, Novartis, Menarini, Sanofi/Regeneron, Roche, Fondazione Maugeri, Fondazione Chiesi, Edmond Pharma. **LP** reports grants, personal fees or non-financial support from Novartis, AstraZeneca, GSK, TEVA, Boehringer-Ingelheim, Chiesi, Sanofi, Menarini, Mundipharma, and Esteve. **MP** and **CR** received funding from EAACI. **SQ** reports personal fees and non-financial support from GSK, AZ, Sanofi, Novartis, Mundipharma, Teva, and Allergy Therapeutics. **JS** declares personal fees from Novartis, GSK, AstraZeneca, and Sanofi. **MS** reports personal fees or research funding from ASIT Biotech.sa, Allergy Therapeutics, ALK, Regeneron, Merck, Immune Tolerance Network, Leti Laboratorios, and Allergopharma. **YS** and **PA-C** has received funding from EAACI. **OP** received research grants from Immunotek S.L. and Novartis; received fees for giving scientific lectures from: Allergy Therapeutics, Amgen, AstraZeneca, Diater, GSK, Immunotek S.L, Novartis, Sanofi-Genzyme and Stallergenes; participated in advisory boards from Novartis and Sanofi-Genzyme. **MK** reports personal fees from ALK-Abello, Allergopharma , Stallergenes, Anergis, Allergy Therapeutics, Circassia, Leti, Biomay, HAL, Astra-Zeneca, GSK, Novartis, Teva, Vectura, UCB, Takeda, Roche, Janssen, MedImmune, and Chiesi. All other authors have no conflict of interest within the scope of the submitted work.

References

1. <http://www.globalasthmareport.org/burden/burden.php>; accessed at 16.12.2019
2. Papi A, Brightling C, Pedersen SE, Reddel HK. Asthma. *Lancet*. 2018;391(10122):783-800.
3. Settipane RA, Kreindler JL, Chung Y, Tkacz J. Evaluating direct costs and productivity losses of patients with asthma receiving GINA 4/5 therapy in the United States. *Ann Allergy Asthma Immunol*. 2019;123(6):564-572
4. Pérez de Llano L, Martínez-Moragón E, Plaza Moral V, Trisan Alonso A, Sánchez CA, Callejas FJ, et al. Unmet therapeutic goals and potential treatable traits in a population of patients with severe uncontrolled asthma in Spain. ENEAS study. *Respir Med*. 2019;151:49-54.
5. Chung KF, Wenzel SE, Brozek JL, Bush A, Castro M, Sterk PJ, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J* 2014; 43: 343–373
6. Corren J, Yawn BP. Advancing the Care of Severe Asthma: Differential Diagnosis, Multidisciplinary Management, and Patient Engagement. *Am J Med*. 2019 Feb 20. pii: S0002-9343(19)30146-9. doi: 10.1016/j.amjmed.2019.02.002.
7. <https://ginasthma.org/wp-content/uploads/2019/06/GINA-2019-main-report-June-2019-wms.pdf>; accessed at 16.12.2019
8. Seys SF, Quirce S, Agache I, Akdis CA, Alvaro-Lozano M, Antolín-Amérigo D, et al. Severe asthma: Entering an era of new concepts and emerging therapies: Highlights of the 4th international severe asthma forum, Madrid, 2018. *Allergy* 2019;74(11):2244-2248
9. Green RH, Brightling CE, McKenna S, Hargadon B, Parker D, Bradding P, et al. Asthma exacerbations and sputum eosinophil counts: a randomised controlled trial. *Lancet*. 2002;360(9347):1715-21.
10. Nair P, O'Byrne PM. Measuring Eosinophils to Make Treatment Decisions in Asthma. *Chest*. 2016;150(3):485-7.
11. Petsky HL, Li A, Chang AB. Tailored interventions based on sputum eosinophils versus clinical symptoms for asthma in children and adults. *Cochrane Database Syst Rev*. 2017;8:CD005603.
12. Stokes JR, Casale TB. Characterization of asthma endotypes: implications for therapy. *Ann Allergy Asthma Immunol*. 2016;117(2):121-5.
13. Agache I. Severe asthma phenotypes and endotypes. *Semin Immunol*. 2019; 46:101301.
14. Agache I, Akdis CA. Precision medicine and phenotypes, endotypes, genotypes, regiotypes, and theratypes of allergic diseases. *J Clin Invest*. 2019;130:1493-1503

15. Agache I, Strasser DS, Klenk A, Agache C, Farine H, Ciobanu C, et al. Serum IL-5 and IL-13 consistently serve as the best predictors for the blood eosinophilia phenotype in adult asthmatics. *Allergy*. 2016;71(8):1192-202.
16. Palomares Ó, Sánchez-Ramón S, Dávila I, Prieto L, Pérez de Llano L, Leonart M, et al. dlvergEnt: How IgE Axis Contributes to the Continuum of Allergic Asthma and Anti-IgE Therapies. *Int J Mol Sci*. 2017;18(6). pii: E1328.
17. <https://www.ema.europa.eu/en/medicines/human/EPAR/nucala>; accessed at 16.12.2019
18. https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761122s000lbl.pdf; accessed at 16.12.2019
19. <https://www.ema.europa.eu/en/medicines/human/EPAR/cinquaero>; accessed at 16.12.2019
20. https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/761033lbl.pdf; accessed at 16.12.2019
21. <https://www.ema.europa.eu/en/medicines/human/EPAR/fasenra>; accessed at 16.12.2019
22. https://www.accessdata.fda.gov/drugsatfda_docs/nda/2017/761070Orig1s000Approv.pdf; accessed at 16.12.2019
23. <https://www.ema.europa.eu/en/medicines/human/summaries-opinion/dupixent-0>; accessed at 16.12.2019
24. https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/761055s007lbl.pdf; accessed at 16.12.2019
25. <https://www.ema.europa.eu/en/medicines/human/EPAR/xolair>; accessed at 16.12.2019
26. https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/103976s5225lbl.pdf; accessed at 16.12.2019
27. Walford HH, Doherty TA. Diagnosis and management of eosinophilic asthma: a US perspective 2014;7:53-65.
28. Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al; Cochrane Bias Methods Group; Cochrane Statistical Methods Group. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928
29. Evers S, Goossens M, de Vet H, van Tulder M, Ament A. Criteria list for assessment of methodological quality of economic evaluations: Consensus on Health Economic Criteria. *Int J Technol Assess Health Care*. 2005;21(2):240-5.
30. Hutter F, Antoñanzas F. Economic evaluations in the EURONHEED: a comparative analysis. *Pharmacoeconomics*. 2009;27(7):561-70.

- Accepted Article
31. Nixon J, Rice S, Drummond M, Boulenger S, Ulmann P, de Pouvourville G. Guidelines for completing the EURONHEED transferability information checklists. *Eur J Health Econ.* 2009;10(2):157-65
 32. Rücker G, Cates CJ, Schwarzer G. Methods for including information from multi-arm trials in pairwise meta-analysis. *Res Synth Methods.* 2017;8(4):392-403.
 33. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, Schünemann HJ; GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ.* 2008;336(7650):924-6.
 34. Guyatt GH, Oxman AD, Sultan S, Glasziou P, Akl EA, Alonso-Coello P, et al; GRADE Working Group. GRADE guidelines. Rating up the quality of evidence. *J Clin Epidemiol.* 2011;64(12):1311-6
 35. Santanello NC, Zhang J, Seidenberg B, Reiss TF, Barber BL. What are minimal important changes for asthma measures in a clinical trial? *Eur Respir J.* 1999;14(1):23-7.
 36. Jones PW. Interpreting thresholds for a clinically significant change in health status in asthma and COPD. *Eur Respir J.* 2002;19(3):398-404
 37. Juniper EF, Svensson K, Mörk AC, Ståhl E. Measurement properties and interpretation of three shortened versions of the asthma control questionnaire. *Respir Med.* 2005;99(5):553-8
 38. Dweik RA, Boggs PB, Erzurum SC, Irvin CG, Leigh MW, Lundberg JO, et al; American Thoracic Society Committee on Interpretation of Exhaled Nitric Oxide Levels (FENO) for Clinical Applications. An official ATS clinical practice guideline: interpretation of exhaled nitric oxide levels (FENO) for clinical applications. *Am J Respir Crit Care Med.* 2011;184(5):602-15.
 39. Bleecker ER, FitzGerald JM, Chanez P, Papi A, Weinstein SF, Barker P, et al; SIROCCO study investigators. Efficacy and safety of benralizumab for patients with severe asthma uncontrolled with high-dosage inhaled corticosteroids and long-acting β_2 -agonists (SIROCCO): a randomised, multicentre, placebo-controlled phase 3 trial. *Lancet.* 2016 ;388(10056):2115-2127
 40. FitzGerald JM, Bleecker ER, Nair P, Korn S, Ohta K, Lommatzsch M, et al; CALIMA study investigators. Benralizumab, an anti-interleukin-5 receptor α monoclonal antibody, as add-on treatment for patients with severe, uncontrolled, eosinophilic asthma (CALIMA): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet.* 2016;388(10056):2128-2141

41. Nair P, Wenzel S, Rabe KF, Bourdin A, Lugogo NL, Kuna P, et al; ZONDA Trial Investigators. Oral Glucocorticoid-Sparing Effect of Benralizumab in Severe Asthma. *N Engl J Med*. 2017;376(25):2448-2458.
42. Wenzel S, Castro M, Corren J, Maspero J, Wang L, Zhang B, et al. Dupilumab efficacy and safety in adults with uncontrolled persistent asthma despite use of medium-to-high-dose inhaled corticosteroids plus a long-acting β 2 agonist: a randomised double-blind placebo-controlled pivotal phase 2b dose-ranging trial. *Lancet*. 2016;388(10039):31-44
43. Castro M, Corren J, Pavord ID, Maspero J, Wenzel S, Rabe KF, et al. Dupilumab Efficacy and Safety in Moderate-to-Severe Uncontrolled Asthma. *N Engl J Med*. 2018;378(26):2486-2496.
44. Rabe KF, Nair P, Brusselle G, Maspero JF, Castro M, Sher L, et al. Efficacy and Safety of Dupilumab in Glucocorticoid-Dependent Severe Asthma. *N Engl J Med*. 2018;378(26):2475-2485.
45. Chupp GL, Bradford ES, Albers FC, Bratton DJ, Wang-Jairaj J, Nelsen LM, et al. Efficacy of mepolizumab add-on therapy on health-related quality of life and markers of asthma control in severe eosinophilic asthma (MUSCA): a randomised, double-blind, placebo-controlled, parallel-group, multicentre, phase 3b trial. *Lancet Respir Med*. 2017;5(5):390-400.
46. Bel EH, Wenzel SE, Thompson PJ, Prazma CM, Keene ON, Yancey SW, et al; SIRIUS Investigators. Oral glucocorticoid-sparing effect of mepolizumab in eosinophilic asthma. *N Engl J Med*. 2014;371(13):1189-97
47. Ortega HG, Liu MC, Pavord ID, Brusselle GG, FitzGerald JM, Chetta A, et al; MENSA Investigators. Mepolizumab treatment in patients with severe eosinophilic asthma. *N Engl J Med*. 2014 ;371(13):1198-207.
48. Hanania NA, Wenzel S, Rosén K, Hsieh HJ, Mosesova S, Choy DF, et al. Exploring the effects of omalizumab in allergic asthma: an analysis of biomarkers in the EXTRA study. *Am J Respir Crit Care Med*. 2013;187(8):804-11.
49. Sorkness CA, Wildfire JJ, Calatroni A, Mitchell HE, Busse WW, O'Connor GT, et al. Reassessment of omalizumab-dosing strategies and pharmacodynamics in inner-city children and adolescents. *J Allergy Clin Immunol Pract*. 2013;1(2):163-71
50. Busse W, Spector S, Rosén K, Wang Y, Alpan O. High eosinophil count: a potential biomarker for assessing successful omalizumab treatment effects. *J Allergy Clin Immunol*. 2013;132(2):485-6

51. Casale TB, Chipps BE, Rosén K, Trzaskoma B, Haselkorn T, Omachi TA, et al. Response to omalizumab using patient enrichment criteria from trials of novel biologics in asthma. *Allergy*. 2018;73(2):490-497.
52. Castro M, Mathur S, Hargreave F, Boulet LP, Xie F, Young J, et al; Res-5-0010 Study Group. Reslizumab for poorly controlled, eosinophilic asthma: a randomized, placebo-controlled study. *Am J Respir Crit Care Med*. 2011;184(10):1125-32
53. Castro M, Zangrilli J, Wechsler ME, Bateman ED, Brusselle GG, Bardin P, et al. Reslizumab for inadequately controlled asthma with elevated blood eosinophil counts: results from two multicentre, parallel, double-blind, randomised, placebo-controlled, phase 3 trials. *Lancet Respir Med*. 2015;3(5):355-66.
54. Corren J, Weinstein S, Janka L, Zangrilli J, Garin M. Phase 3 Study of Reslizumab in Patients With Poorly Controlled Asthma: Effects Across a Broad Range of Eosinophil Counts. *Chest*. 2016;150(4):799-810
55. Bjermer L, Lemiere C, Maspero J, Weiss S, Zangrilli J, Germinaro M. Reslizumab for Inadequately Controlled Asthma With Elevated Blood Eosinophil Levels: A Randomized Phase 3 Study. *Chest*. 2016;150(4):789-798
56. https://icer-review.org/wpcontent/uploads/2018/04/ ICER_Asthma_Final_Report_122018-1.pdf; accessed at 18.12.2019
57. Tikhonova I, Long L, Ocean N, Barnish M, Robinson S, Nikram S, et al. Benralizumab for treating severe eosinophilic asthma: NICE single technology appraisal. *Value in health* 2019; 22; suppl.2:358
58. Whittington MD, McQueen RB, Ollendorf DA, Tice JA, Chapman RH, Pearson SD, et al. Assessing the value of mepolizumab for severe eosinophilic asthma: a cost-effectiveness analysis. *Ann Allergy Asthma Immunol*. 2017;118(2):220-225
59. Han S, Kim S, Kim H, Suh HS. Cost-utility analysis of reslizumab for patients with severe eosinophilic asthma inadequately controlled with high-dose inhaled corticosteroids and long-acting beta2-agonists in South Korea. *Current medical research and opinion*. 2019; 10.1080/03007995.2019.1605159. [Epub ahead of print]
60. Lam J, Hay JW, Salcedo J, Kenyon NJ. A cost-effectiveness analysis of reslizumab in the treatment of poorly controlled eosinophilic asthma. *J Asthma*. 2019; 56 (8): 872-881
61. Liu W, Ma X, Zhu W. Adverse events of benralizumab in moderate to severe eosinophilic asthma. A meta-analysis. *Medicine (Baltimore)*. 2019;98(22):e15868.
62. Tian BP, Zhang GS, Lou J, Zhou HB, Cui W. Efficacy and safety of benralizumab for eosinophilic asthma: A systematic review and meta-analysis of randomized controlled trials. *J Asthma*. 2018; 55(9):956-965

63. Bassani C, Rossi L, Siveris K, Sferelli RL, Saraiva L, Tanno LK.. Use of dupilumab on the treatment of moderate-to-severe asthma: a systematic review. *Rev Assoc Med Bras* (1992). 2019;65(9):1223-1228
64. Yancey SW, Ortega HG, Keene ON, Mayer B, Gunsoy NB, Brightling CE, et al. Meta-analysis of asthma-related hospitalization in mepolizumab studies of severe eosinophilic asthma. *J Allergy Clin Immunol*. 2017;139(4):1167-1175
65. Powell C, Milan SJ, Dwan K, Bax L, Walters N. Mepolizumab versus placebo for asthma. *Cochrane Database Syst Rev*. 2015;(7):CD010834
66. Colombo GL, Di Matteo S, Martinotti C, Oselin M, Valentino MC, Bruno GM, et al. Omalizumab and long-term quality of life outcomes in patients with moderate-to-severe allergic asthma: a systematic review. *Ther Adv Respir Dis*. 2019;13:1753466619841350
67. Corren J, Kavati A, Ortiz B, Vegesna A, Colby JA, Ruiz K, et al. Patient-reported outcomes in moderate-to-severe allergic asthmatics treated with omalizumab: a systematic literature review of randomized controlled trials. *Curr Med Res Opin*. 2018; 34(1):65-80
68. Li J, Wang F, Lin C, Du J, Xiao B, Du C, Sun J. The efficacy and safety of reslizumab for inadequately controlled asthma with elevated blood eosinophil counts: A systematic review and meta-analysis. *J Asthma*. 2017;54(3):300-307
69. McQueen RB, Sheehan DN, Whittington MD, van Boven JFM, Campbell JD. Cost-Effectiveness of Biological Asthma Treatments: A Systematic Review and Recommendations for Future Economic Evaluations. *Pharmacoeconomics*. 2018;36(8):957-971
70. Lundh A, Lexchin J, Mintzes B, Schroll JB, Bero L. Industry sponsorship and research outcome. *Cochrane Database Syst Rev*. 2017;2:MR000033.
71. Goldman M, Hirsch I, Zangrilli JG, Newbold P, Xu X. The association between blood eosinophil count and benralizumab efficacy for patients with severe, uncontrolled asthma: subanalyses of the Phase III SIROCCO and CALIMA studies. *Current Medical Research and Opinion* 2017; 3(9):1605-1613
72. Aburuz S, McElnay J, Gamble J, Millership J, Heaney L. Relationship between lung function and asthma symptoms in patients with difficult to control asthma. *J Asthma*. 2005;42(10):859-64.
73. Wagener AH, de Nijs SB, Lutter R, Sousa AR, Weersink EJ, Bel EH, et al. External validation of blood eosinophils, FE(NO) and serum periostin as surrogates for sputum eosinophils in asthma. *Thorax* 2015, 70(2):115-20

74. Hastie A, Moore W, Li H, Rector B, Ortega V, Pascual R, et al. Biomarker Surrogates Do Not Accurately Predict Sputum Eosinophils and Neutrophils in Asthma. *J Allergy Clin Immunol* 2013; 132(1):72-80.
75. Strauss RA, Jawhari N. Mepolizumab in the treatment of severe eosinophilic asthma: Results from a physician in the field. *Ann Allergy Asthma Immunol*. 2018;121(1):121-123
76. Pelaia C, Busceti MT, Solinas S, Terracciano R, Pelaia G. Real-life evaluation of the clinical, functional, and hematological effects of mepolizumab in patients with severe eosinophilic asthma: Results of a single-centre observational study. *Pulm Pharmacol Ther*. 2018;53:1-5.
77. Ortega H, Menzies-Gow A, Llanos JP, Forshag M, Albers F, Gunsoy N, et al. Rapid and Consistent Improvements in Morning PEF in Patients with Severe Eosinophilic Asthma Treated with Mepolizumab. *Adv Ther*. 2018;35(7):1059-1068.
78. Pelaia C, Calabrese C, Barbuto S, Busceti MT, Preianò M, Gallelli L, et al. Omalizumab lowers asthma exacerbations, oral corticosteroid intake and blood eosinophils: Results of a 5-YEAR single-centre observational study. *Pulm Pharmacol Ther*. 2019;54:25-30.
79. Bime C, Wei CY, Holbrook JT, Sockrider MM, Revicki DA, Wise RA. . Asthma Symptom Utility Index: Reliability, validity, responsiveness and the minimal important difference in adult asthma patients. *J Allergy Clin Immunol*. 2012; 130(5):1078-84
80. Brussino, L, Heffler, E, Bucca, C, Nicola, S, Rolla, G. Eosinophils Target Therapy for Severe Asthma: Critical Points. *BioMed Research International*; 2018. 7582057

Figure 1. Study flowchart for the selecting evidence for efficacy and safety of the biologicals in severe eosinophilic asthma




Figure 2. Study flowchart for selecting the economic evidence for biologicals in severe eosinophilic asthma




Table 1. Summary of the characteristics of the included studies evaluating clinical efficacy of benralizumab, mepolizumab, omalizumab and reslizumab as add-on for uncontrolled severe asthma

Author, year (Trial Name)	Outcomes assessment	Age (years) Mean \pm SD Range	Population (N)	Intervention Type Number of patients exposed (N)	Control
Benralizumab					
Bleecker 2016 (39) (SIROCCO)	48	47.6 \pm 14.5 12-75	Severe, uncontrolled asthma (N=1205)	Benralizumab: 30 mg SC Q8W (N=398) Benralizumab: 30 mg SC Q4W (N=400)	Standard of care (SOC)
FitzGerald 2016 (40) (CALIMA)	56	49.0 \pm 14.3 12-75	Severe, uncontrolled asthma (N= 1306)	Benralizumab: 30 mg SC Q8W (N=441) Benralizumab 30mg SC Q4W (N=425)	SOC
Nair 2017 (41) (ZONDA)	24	52.9 \pm 10.1 18-75	Severe uncontrolled asthma receiving daily OCS (N=220)	Benralizumab 30mg SC Q8W (N=73) Benralizumab 30mg SC Q4W (N=72)	SOC
Dupilumab					
Castro 2018 (43) (Liberty Asthma QUEST)	52	47.9 \pm 15.3 >12	Uncontrolled asthma (N=1902)	Dupilumab 200 mg SC Q2W (loading dose, 400mg) (N=631) Dupilumab 300 mg SC Q2W (loading dose, 600mg) (N=633)	SOC
Rabe 2018 (44)	24	51.3 \pm 12.6 >18	Oral glucocorticoid-treated asthma (N=210)	Dupilumab 300 mg SC Q2W (N=103)	SOC
Wenzel 2016 (42)	24	48.0 \pm 12,8 >18	Moderate to severe, uncontrolled asthma (N=776)	Dupilumab 200 mg SC Q2W Dupilumab 300 mg SC Q2W Dupilumab 200 mg SC Q4W Dupilumab 300 mg SC Q4W	SOC
Mepolizumab					

Bel 2014 (46) (SIRIUS)	24	50 16-74	Severe eosinophilic asthma receiving daily OCS (N=135)	100 mg SC Q4W (N=69)	SOC
Ortega 2014 (47) (MENSA)	32	51 12-81	Severe eosinophilic asthma (N=576)	75-mg IV Q4W (N=191) 100-mg SC Q4W (N=194)	SOC
Chupp 2017 (45) (MUSCA)	24	49.8 >12	Severe eosinophilic asthma (N=551)	100 mg SC Q4W (N=274)	SOC
Omalizumab					
Casale 2018 (51)	16	39.7 ± 13.8 12-75	Eosinophilic asthma (N=1071)	SC Q2W or Q4W at a dose calculated based on body weight and baseline serum IgE levels (150 mg or 300 mg Q4W or 225 mg, 300 mg, or 375 mg Q2W) (N=542)	SOC
Hanania 2013 (48)	48	44 ± 15 12-75	Uncontrolled severe persistent allergic asthma (N=850) Subgroup high blood eosinophils (N=414) or high FeNO (N=101)	Dose determined by the European omalizumab dosing table, ensuring a minimum omalizumab dose of 0.008 mg/kg/IgE (IU/mL) Q2W or a minimum of 0.016 mg/kg/IgE (IU/mL) Q4W.	SOC
Sorkness 2013 (49)	48	10.7 ± 3.6 6-20	Moderate to severe persistent allergic asthma Subgroup with high FeNO (N=155)	Dose determined the expanded FDA dosage table, ensuring a minimum dose of 0.016 mg/kg/IgE (IU/mL) every month	SOC
Busse 2013 (50)	28	12-75	Patients with atopic asthma Subgroup with eosinophilic asthma (N=51)	Dose determined by the European omalizumab dosing table, ensuring a minimum omalizumab dose of 0.008 mg/kg/IgE (IU/mL) Q2W or a minimum of 0.016 mg/kg/IgE (IU/mL) Q4W	SOC
Reslizumab					

Bjermer 2016 (55)	16	43 (12-75)	Inadequately controlled eosinophilic asthma (N=311)	Reslizumab 3.0 mg/kg IV Q4W (N=106)	SOC
Castro 2011 (52)	12	44.9 ± 13.94 (18-75)	Inadequately controlled eosinophilic asthma (N=106)	Reslizumab 3.0 mg/kg IV Q4W (N=53)	SOC
Castro 2015 (53)	52	49 (12-75)	Inadequately controlled eosinophilic asthma (N=953)	Reslizumab 3 mg/kg IV Q4W (N=477)	SOC
Corren 2016 (54)	16	44.9 (18-65)	Inadequately controlled asthma Subgroup with baseline eosinophils ≥ 400 cells/μL (n=96)	Reslizumab 3 mg/kg IV Q4W (N=77)	SOC

Table 2. Summary of the characteristics of the included studies evaluating the economical impact of biologicals as add-on for uncontrolled severe asthma

Author, year	Design, Country	Intervention	Control	Time horizon, perspective	Difference in cost (year value)	Difference in outcome	ICER	Risk of bias (CHEC score)	Transferability score	Source of funding
Benralizumab										
ICER 2019 (56)	Cost-utility Markov model, US	Lifetime Benralizumab in addition to SOC	Lifetime SOC	Lifetime, US health care perspective. Societal in the sensitivity analysis.	581,000 \$ (2018 US Dollars)	1.41 QALY	412,000 \$ / QALY	17.5/20	13.5/16	Government grants and non-profit foundations
Tikhonova 2018 (57)	Cost-utility Markov model, UK	Lifetime Benralizumab in addition to SOC	Lifetime SOC	Lifetime, UK NHS perspective. Societal in the sensitivity analysis.	Not Reported	QALY not reported	39,135 £ / QALY	15.5/20	12.5/16	AstraZeneca financed the original model critiqued by the University of Exeter
Dupilumab										
ICER 2019 (56)	Cost-utility Markov model, US	Lifetime Dupilumab in addition to SOC	Lifetime SOC	Lifetime, US health care perspective. Societal in the sensitivity analysis.	704,000 \$ (2018 US Dollars)	1.51 QALY	464,000 \$ / QALY	17.5/20	13.5/16	Government grants and non-profit foundations
Mepolizumab										
Whittington 2017 (58)	Cost-utility Markov model, US	Lifetime Mepolizumab in addition to SOC	Lifetime SOC	Lifetime, health system	589,941 \$ (2014 US Dollar)	1.53 QALY	385,546 \$ / QALY	16/20	14/16	Institute for Clinical and Economic Review

Author, year	Design, Country	Intervention	Control	Time horizon, perspective	Difference in cost (year value)	Difference in outcome	ICER	Risk of bias (CHEC score)	Transferability score	Source of funding
Reslizumab										
Han 2019 (59)	Cost-utility Markov model, South Korea	Lifetime Reslizumab in addition to SOC	Lifetime SOC	Lifetime, societal	119,394 \$ (2018 US Dollars)	5.17 QALY	23,081 \$ / QALY	16.5/20	13/16	Teva-Handok Pharma
Lam 2018 (60)	Cost-utility Markov model, US	5-year Reslizumab in addition to SOC	5-year SOC	5-year, societal	24,404 \$ (2017 US Dollar)	0.035 QALY	697,403 \$ / QALY	16/20	12.5/16	University of Southern California, US

Table 3: Summary of findings for Benralizumab compared to standard of care for eosinophilic asthma					
Outcomes	No of participants (studies) Follow-up (range)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with standard of care	Risk difference with benralizumab
Exacerbations Assessed with annualised asthma exacerbation rate	1373 (3 RCTs) ^{39,40,41} 28 to 56 weeks	⊕⊕⊕⊕ HIGH ^{3,a,b}	Rate ratio 0.53 (0.39 to 0.72) ^{c,d}	1500 exacerbations per 1000 patients per year	705 fewer exacerbations per 1.000 patients per year (915 fewer to 420 fewer)
Asthma Control Assessed with ACQ-6 score between-group-difference at the end of the study	1373 (3 RCTs) ^{39,40,41} 28 to 56 weeks	⊕⊕⊕⊕ HIGH ^{3,4,b,e,f}	-		mean difference - 0.26 (- 0.46 to - 0.07 fewer) ^{c,g}
Quality of life Assessed with Asthma Quality of Life Questionnaire for 12 years and older	1333 (3 RCTs) ^{39,40,41} 28 to 52 weeks	⊕⊕⊕⊕ HIGH ^{3,6,b,j,k}	-		mean difference+ 0.23 higher (+0.11 to +0.36) ^c
Any drug related adverse event (AE) Assessed with number of events	478 (1 RCT) ⁴⁰ 56 weeks	⊕⊕⊕○ MODERATE ^{3,b,l}	Rate ratio 1.41 (0.87 to 2.27)	105 per 1.000	43 more per 1.000 (14 fewer to 133 more)
Any serious adverse event (SAE) unrelated to asthma exacerbation Assessed with number of events	148 (1 RCT) ⁴¹ 28 weeks	⊕⊕○○ LOW ^{3,b,l}	Rate ratio 0.56 (0.22 to 1.44)	147 per 1.000	65 fewer per 1.000 (114 fewer to 65 more)
Decrease in OCS use Assessed with reduction in daily OCS dose of ≥50%	148 (1 RCT) ⁴¹ 28 weeks	⊕⊕⊕⊕ HIGH ^{3,b}	Rate ratio 1.76 (1.26 to 2.47)	373 per 1.000	284 more per 1.000 (97 more to 549 more)
Lung function Assessed with pre-bronchodilator FEV1 (mL) between-group-difference at the end of the study	1370 (3 RCTs) ^{39,40,41} 28 to 56 weeks	⊕⊕⊕○ MODERATE ^{3,4,5,b,h,i}	-		mean difference + 140 mL (+90 to +190 higher) ^c
Rescue medication use Assessed with puffs/day	0 studies	-	not estimable		

Table 3: Summary of findings for Benralizumab compared to standard of care for eosinophilic asthma

Outcomes	No of participants (studies) Follow-up (range)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with standard of care	Risk difference with benralizumab

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

GRADE Working Group grades of evidence

High certainty: High confidence that the true effect lies close to that of the estimate of the effect

Moderate certainty: Moderate confidence in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Limited confidence in the effect estimate: The true effect may be substantially different from the estimate of the effect

Very low certainty: Little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations

a. Statistically significant ($I^2=65\%$) but probably unimportant heterogeneity.

b. All included studies were funded by industry, and all showed positive results. No industry-independent observational or randomized studies were identified to contrast results. Therefore, the quality of the evidence was downgraded for potential publication bias (70).

c. The pooled data were assessed at 28 weeks (41) and at 48-52 weeks (71). Goldman 2017 included patients aged 12-17 years old.

d. In the current systematic review 2 studies reporting the effect on exacerbation leading to emergency room visits or hospitalizations were also included. The pooled risk ratio was 0.24 (95% CI 0.03-1.72) (see full text report).

e. Statistically significant ($I^2=61\%$) but probably unimportant heterogeneity.

f. The minimal important difference (MID) for ACQ-6 is 0.5 points (35).

g. In the current systematic review 3 studies reporting the effect on total asthma control score change were also included. The pooled mean difference was -0.19 (95% CI -0.31 to -0.08), see full text report.

h. Quality of the evidence was downgraded because FEV1 is considered a surrogate outcome for asthma control, with a variable correlation with asthma symptoms (72).

i. The panel agreed that minimal important difference for FEV1 is 0.20 L.

j. Statistically significant ($I^2=55\%$) but probably unimportant heterogeneity.

k. For AQLQ(S)+12 the MID is 0.5 (37)

l. The effect may both be harmful or beneficial. Small sample size and number of events.

Table 4: Summary of findings of Dupilumab compared to standard of care for eosinophilic asthma

Outcomes	No of participants (studies) Follow-up (range)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with standard of care	Risk difference with dupilumab
Exacerbations Assessed with annualised asthma exacerbation rate	1712 (3 RCTs) ^{42,43,44} 24 to 52 weeks	⊕⊕⊕⊕ HIGH ^{4,a,b}	Rate ratio 0.44 (0.32 to 0.59)	1570 exacerbations per 1000 patients per year	894 fewer exacerbations per 1000 patients per year (1086 fewer to 655 fewer) ^c
Asthma control assessed with: Asthma Control Questionnaire -5 Scale from: 1 to 5	507 (1 RCT) ⁴² 24 weeks	⊕⊕⊕○ MODERATE ^{4,8,a,b,g}	-		mean difference - 0.48 (-0.88 lower to - 0.09)
Quality of life Assessed with asthma Quality of Life Questionnaire Scale from: 1 to 7	958 (2 RCTs) ^{43,44} 24 to 52 weeks	⊕⊕⊕○ MODERATE ^{4,9,a,b,h}	-		mean difference + 0.42 higher (+0.25 to +0.59)
Treatment-related adverse events (AE) Assessed with number of events	264 (1 RCT) ⁴² 24 weeks	⊕⊕⊕○ MODERATE ^{4,a,b,m}	RR 1.00 (0.88 to 1.13)	794 per 1.000	0 fewer per 1.000 (95 fewer to 103 more)
Treatment-related serious adverse events (SAE) Assessed with number of events	264 (1 RCT) ⁴² 24 weeks	⊕⊕○○ LOW ^{4,a,b,m}	RR 1.46 (0.60 to 3.54)	59 per 1.000	27 more per 1.000 (24 fewer to 149 more)
Decrease in OCS dose Assessed with percentage of reduction compared to baseline	150 (1 RCT) ⁴² 24 weeks	⊕⊕⊕⊕ HIGH ^{4,a,b}	-		mean difference - 29.4 % (-43.23 to -15.57)
Lung function Assessed with FEV1 in mL	1030 (3 RCTs) ^{42,43,44} 24 to 52 weeks	⊕⊕○○ LOW ^{4,5,6,7,a,b,d,e,f}	-		mean difference + 180 mL (+110 to +250)
Fraction of exhaled nitric oxide Assessed with mean % change (ppb) from baseline	150 (1 RCT) ⁴² 24 weeks	⊕⊕○○ LOW ^{4,10,11,12,a,b,i,j}	-		mean difference - 40.11 % (-78.68 to -1.55)
Rescue medication use Assessed with puffs/day	143 (1 RCT) ⁴² 24 to 52 weeks	⊕⊕⊕○ MODERATE ^{4,7,a,b,k,l}	-		mean difference - 0.56 puff/day (-2.28 to +1.16)

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

Table 4: Summary of findings of Dupilumab compared to standard of care for eosinophilic asthma

Outcomes	No of participants (studies) Follow-up (range)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with standard of care	Risk difference with dupilumab

GRADE Working Group grades of evidence

High certainty: High confidence that the true effect lies close to that of the estimate of the effect

Moderate certainty: Moderate confidence in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Limited confidence in the effect estimate: The true effect may be substantially different from the estimate of the effect

Very low certainty: Little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations

a. All included studies had a high risk of selective reporting bias (42,43,44). However, the evidence quality was not downgraded because most of the outcomes of interest for our analysis were reported.

b. All included studies were founded by industry and the same company (Sanofi and Regeneron Pharmaceuticals), and all showed positive results. No industry-independent observational or randomized trials were identified to contrast the results. Therefore, the quality of the evidence was downgraded for potential publication bias (70)

c. Two studies (Rabe 2018, Wenzel 2016) assessed exacerbations at 24 weeks, and Castro2018 at 52 weeks.

d. The quality of the evidence was downgraded because FEV1 is considered a surrogate outcome of asthma control, with a variable correlation with asthma symptoms (72).

e. The panel agreed that minimal important difference (MID) for FEV1 is 0.20 L and considered the effect as imprecise.

f. The panel agreed that minimal important difference (MID) for FEV1 is 0.20 L and thus the effect was considered as imprecise.

g. Downgraded because the effect of dupilumab is beneficial but the lower side of the CI is less than the MID(0.5 points). (37)

h. Downgraded because the effect of dupilumab is beneficial but the lower side of the CI is less than the MID(0.5 points). (37)

i. Downgraded because FeNO is not consistently considered a good surrogate of eosinophilic inflammation (73,74)

j. From one visit to the next a change greater than 20% for basal values over 50 ppb or more than 10 ppb for basal values lower than 50 ppb may indicate significant response (38).

k. Downgraded because the effect may both be beneficial and harmful.

l. The MID for rescue medication use is a reduction by 0.81 puffs/day (35).

m. The effect may both be harmful or beneficial. Small number of events.

Table 5: Summary of findings of mepolizumab compared to standard of care for eosinophilic asthma

Outcomes	No of participants (studies) Follow-up (range)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with standard of care	Risk difference with mepolizumab
Exacerbations Exacerbation rate ratio Assessed with the annualized rates of asthma exacerbations	1071 (3 RCTs) ^{45,46,47} 24 to 32 weeks	⊕⊕⊕⊕ HIGH ^{4,5,a,b,c}	Rate ratio 0.49 (0.38 to 0.66)	1700 exacerbations per 1000 patients per year	870 fewer exacerbations per 1000 patients per year (592 fewer to 1079 fewer)
Exacerbations leading to hospitalisation Assessed with the annualized rate of asthma exacerbations leading to hospitalisation	(2 RCTs) ^{45,47} 24 to 32 weeks	⊕⊕⊕⊕ HIGH ^{4,5,a,c,d}	Rate ratio 0.30 (0.13 to 0.71)	100 exacerbations per 1000 patients per year	70 fewer exacerbations per 1000 patients per year (29 fewer to 87 fewer)
Asthma control Assessed with: ACQ-5 score between-group-difference at the end of the study Scale from: 0 to 6 ^{9,j}	912 (3 RCTs) ^{45,47}	⊕⊕⊕○ MODERATE ^{4,5,a,c,i}	-		mean difference - 0.43 (- 0.56 to - 0.31)
Quality of life Assessed with St. George's Respiratory Questionnaire between-group-difference at the end of the study	1045 (3 RCTs) ^{45,46,47} 24 to 32 weeks ^{10,k}	⊕⊕⊕○ MODERATE ^{4,5,a,c,i}	-		mean difference - 7.14 (- 9.07 to - 5.21)
Treatment-related adverse events (AE) Assessed with number of events	1071 (3 RCTs) ^{45,46,47}	⊕⊕⊕⊕ HIGH ^{4,5,c}	Rate ratio 1.35 (1.01 to 1.80)	796 per 1.000	279 more per 1.000 (8 more to 637 more)
Treatment-related serious adverse events (SAE) Assessed with number of events	385 (1 RCT) ⁴⁷	⊕○○○ VERY LOW ^{4,5,c,m,n}	Rate ratio 0.98 (0.06 to 15.63)	5 per 1.000	0 fewer per 1.000 (-5 fewer to 77 more)
Lung function assessed with pre-bronchodilator FEV1 (mL) between-group-difference at the end of the study	1043 (3 RCTs) ^{45,46,47} 24 to 32 weeks ^{6,e}	⊕⊕⊕○ MODERATE ^{4,5,7,a,c,f}	-		mean difference +110.9 mL (+58.91 to +162.89)
Lung function assessed with AM peak expiratory flow (PEF)	936 (2 RCTs) ⁷⁷ 24 weeks ^{6,g}	⊕⊕○○ LOW ^{4,5,c,h,i}	-		mean difference +22.46 (+13.98 to +30.94)

Table 5: Summary of findings of mepolizumab compared to standard of care for eosinophilic asthma

Outcomes	No of participants (studies) Follow-up (range)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with standard of care	Risk difference with mepolizumab
Rescue medication use assessed with puffs/day	(1 RCT) ⁴⁵ 21 to 24 weeks ^o	⊕⊕⊕⊕ HIGH ^{4,5,c}	-		mean difference -0.1 puff/day (-0.35 to +0.15)

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

GRADE Working Group grades of evidence

High certainty: High confidence that the true effect lies close to that of the estimate of the effect

Moderate certainty: Moderate confidence in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Limited confidence in the effect estimate: The true effect may be substantially different from the estimate of the effect

Very low certainty: Little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations

a. Two of three studies had a high risk of attrition bias (45,47). Modified intention-to-treat analysis was conducted (i.e. patients were analysed as treated, not as randomized).

b. Probable unimportant heterogeneity

c. Included studies were all funded by industry, and all showed positive results. We identified two industry-independent observational trials that showed similar effects with our meta-analysis (76,77).

d. Mean rates of exacerbation requiring hospitalization across studies were very low (i.e. from 0.02 to 0.10 exacerbations requiring hospitalization per person-year), both in the placebo and intervention arms

e. The panel agreed that minimal important difference (MID) for FEV1 is 0.20 L

f. Downgraded because FEV1 is considered a surrogate outcome of asthma control of symptoms, with a variable correlation with asthma symptoms (72).

g. The MID of PEF is 18.8 L/min (35)

h. Potential attrition bias because PEF baseline values reported in the primary publication (47) differed from values reported in post-hoc analysis publication (77)

i. Downgraded because the lower CI boundary crosses the MID threshold

j. 0.5 points is the minimal important difference for the Asthma Control Questionnaire (ACQ-5 score) (37)

k. >-4.0 was considered the threshold for the MID for quality of life measured with the St.George's Respiratory Questionnaire (36)

l. The St.George's Respiratory Questionnaire SGRQ is not a disease-specific questionnaire for asthma

m. Findings from only 1 RCT available. Downgraded due to publication bias

n. Very few numbers of events per arm

o. The minimal important difference for rescue medication use is -0.81 puffs/day (35)

Table 6: Summary of findings of Omalizumab compared to standard of care for eosinophilic asthma

Outcomes	No of participants (studies) Follow-up (range)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with standard of care	Risk difference with omalizumab
Exacerbations Assessed with annual asthma exacerbations rate	779 (3 RCTs) ^{48,50,51} 16 to 48 weeks	⊕⊕⊕⊕ HIGH ^{4,a,b}	Rate ratio 0.56 (0.40 to 0.77)	660 exacerbations per 1000 patients per year	290 fewer exacerbations per 1.000 patients per year (396 fewer to 152 fewer)
Asthma Control Assessed with Total Asthma Symptoms Score	414 (1 RCT) ⁴⁸ 48 weeks	⊕⊕○○ LOW ^{4,a,b,c,d}	-		mean difference - 0.16 (- 0.51 to +0.19) ^{e,f}
Quality of Life Assessed with Asthma Quality of Life Questionnaire	414 (1 RCT) ⁴⁸ 48 weeks	⊕⊕⊕○ MODERATE ^{4,a,b,d}	-		mean difference + 0.13 (-0.11 to + 0.37) ⁱ
Any adverse event Assessed with number of events	414 (1 RCT) ⁴⁸ 48 weeks	⊕⊕⊕○ MODERATE ^{4,a,b,d}	Rate ratio 1.01 (0.91 to 1.11)	794 per 1.000	8 more per 1.000 (71 fewer to 87 more)
Lung Function Assessed with % pre-bronchodilator FEV1 between-group-difference at the end of the study	(2 RCTs) ^{48,50} 24 weeks to 48 weeks	⊕⊕⊕○ MODERATE ^{4,5,a,b,g,h,i}	-		mean difference + 3.7 % (-2.1 lower to + 9.5) ^{j,k}
Rescue medication use Assessed with puffs/day change from baseline	414 (1 RCT) ⁴⁸ 48 weeks	⊕⊕⊕○ MODERATE ^{4,a,b,d}	-		mean difference - 0.34 (-0.83 to +0.15) ^{6,m,n}

GRADE Working Group grades of evidence**High certainty:** High confidence that the true effect lies close to that of the estimate of the effect**Moderate certainty:** Moderate confidence in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different**Low certainty:** Limited confidence in the effect estimate: The true effect may be substantially different from the estimate of the effect**Very low certainty:** Little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect**Explanations**

a. Information of included studies from non-predefined subgroup analysis.

b. Included studies were all funded by industry, and all showed positive results. We identified one industry-independent observational trial that showed similar effects with our meta-analysis (78).

c. The total asthma symptoms score is an unvalidated scale.

d. The effect may both be harmful or beneficial.

e. Data from subgroup of patients with blood eosinophil count $\geq 260/\mu\text{l}$. This study also reported total asthma symptoms score for the subgroup of FeNO $\geq 24\text{ppb}$, the mean difference is -0.25 (CI 95% -0.77 to 0.27).

- f. In the current systematic review we also included one study (49) reporting the effect on the symptom days over the previous 2 weeks at 48 weeks follow up, the mean difference is -0.45 (p=0.05) (see full text report).
- g. Statistically significant (I²=70%), but probably unimportant heterogeneity.
- h. Downgraded because FEV1 is considered a surrogate outcome of asthma control of symptoms, with a variable correlation with asthma symptoms (72)
- i. The panel agreed that minimal important difference (MID) for FEV1 is 0.20 L j. Population with different threshold of eosinophil counts across the studies: ≥ 200 cells/ μ l (51), ≥ 300 cells/ μ l (50) and ≥ 260 cells/ μ l (48).
- k. One of the included studies (48) also reported the effect on FEV1 % change for a population with FeNO ≥ 24 ppb, the LS mean difference is 3.20 (CI 95% -0.74 to 0.27). The pooled effect evaluated at 48 weeks (48) and 24 weeks (50).
- l. Data from subgroup of patients with blood eosinophil count ≥ 260 /ul. This study also reported AQLQ for the subgroup of FeNO ≥ 24 ppb , the mean difference is 0.37 (CI 95% 0.01 to 0.73).
- m. This study also reported the effect on rescue medication use for the subgroup of FeNO ≥ 24 ppb, the mean difference is -0.49 (CI 95% -0.88 to -0.11).
- n. The MID for rescue medication use is 0.81 puffs/day (35).

References

1. Busse W, Spector S, Rosén K, Wang Y, Alpan O. High eosinophil count: a potential biomarker for assessing successful omalizumab treatment effects. J Allergy Clin Immunol 2013
2. Hanania, NA, Wenzel, S, Rosen, K, Hsieh, HJ, Mosesova, S, Choy, DF, Lal, P, Arron, JR, Harris, JM, Busse, W. Exploring the effects of omalizumab in allergic asthma: An analysis of biomarkers in the EXTRA study. American Journal of Respiratory and Critical Care Medicine; 2013.
3. Casale, TB, Chipps, BE, Rosen, K, Trzaskoma, B, Haselkorn, T, Omachi, TA, Greenberg, S, Hanania, NA. Response to omalizumab using patient enrichment criteria from trials of novel biologics in asthma. Allergy; 2018.
4. Pelaia, C, Busceti, MT, Solinas, S, Terracciano, R, Pelaia, G. Real-life evaluation of the clinical, functional, and hematological effects of mepolizumab in patients with severe eosinophilic asthma: Results of a single-centre observational study. Pulmonary pharmacology & therapeutics; 2018.
5. Aburuz, . Relationship between lung function and asthma symptoms in patients with difficult to control asthma. 2005.

Table 7: Summary of findings of Reslizumab compared to standard of care for eosinophilic asthma

Outcomes	No of participants (studies) Follow-up (mean or range)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with standard of care	Risk difference with reslizumab
Exacerbations Assessed with annualised rate of asthma exacerbations ^a	1059 (3 RCTs) ^{1,2} 52 weeks	⊕⊕⊕⊕ HIGH ^{3,b}	Rate ratio 0.46 (0.37 to 0.58) ^c	1800 exacerbations per 1000 patients per year	972 fewer exacerbations per 1000 patients per year (1134 fewer to 756 fewer)
Exacerbations leading to ER visit or hospitalisation Assessed with annualised rate of asthma exacerbations	953 (2 RCTs) ¹ 52 weeks	⊕⊕⊕○ MODERATE ^{3,b,d}	Rate ratio 0.67 (0.39 to 1.17)	120 exacerbations per 1000 patients per year	40 fewer exacerbations per 1000 patients per year (73 fewer to 20 more)
Asthma control Assessed with Asthma Control Questionnaire-7 Scale from: 0 to 6	1359 (5 RCTs) ^{1,2,4,5} 15 weeks to 16 weeks	⊕⊕⊕⊕ HIGH ^{3,8,b,h}	-		mean difference - 0.25 (- 0.34 to - 0.16)
Quality of life assessed with Asthma Quality of Life Questionnaire Scale from: 1 to 7	1153 (3 RCTs) ^{1,4} 15 to 16 weeks	⊕⊕⊕⊕ HIGH ^{3,10,b,k}	-		mean difference + 0.17 (+0.08 to + 0.25) ^l
Treatment-related adverse events Assessed with number of events	1269 (4 RCTs) ^{1,2,4} 15 to 52 weeks	⊕⊕⊕○ MODERATE ^{3,b,o}	Rate ratio 1.18 (0.89 to 1.56) ^p	125 per 1.000	22 more per 1.000 (14 fewer to 70 more) ^p
Treatment-related serious adverse events Assessed with number of events	1269 (4 RCTs) ^{1,2,4} 15 to 52 weeks ^p	⊕⊕○○ LOW ^{3,b,q}	Rate ratio 4.71 (0.54 to 41.31)	0 per 1.000	0 fewer per 1.000 (0 fewer to 0 fewer)
Decrease in inhaled corticosteroid (ICS) and oral corticosteroid (OCS) dose	0 studies	-	-	-	-
Lung function Assessed with: FEV ₁ in mL	1360 (5 RCTs) ^{1,2,4,5} 15 to 16 weeks	⊕⊕⊕○ MODERATE ^{3,6,7,b,e,f}	-		mean difference + 141.82 mL (+89.23 to 194.41) ⁹⁺
Rescue medication use Assessed with puffs/day	1251 (4 RCTs) ^{1,4,5} 16 weeks	⊕⊕⊕⊕ HIGH ^{3,7,b,n}	-		mean difference - 0.24 (-0.46 to - 0.02)

Table 7: Summary of findings of Reslizumab compared to standard of care for eosinophilic asthma

Outcomes	No of participants (studies) Follow-up (mean or range)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with standard of care	Risk difference with reslizumab
Asthma symptoms Assessed with: Asthma Symptom Utility Index Scale from: 0 to 1	1157 (3 RCTs) ^{1,4} 16 weeks to 16 weeks	⊕⊕⊕⊕ HIGH ^{3,9,b,j}	-		mean difference + 0.05 (+0.03 to +0.07 higher)
Changes in blood eosinophil counts Assessed with: cells/μL	1264 (4 RCTs) ^{1,2,4} 15 weeks to 16 weeks	⊕⊕⊕○ MODERATE ^{3,11,b,m}	-		mean difference - 468.58 (-494.92 to - 442.24)

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

GRADE Working Group grades of evidence

High certainty: High confidence that the true effect lies close to that of the estimate of the effect

Moderate certainty: Moderate confidence in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Limited confidence in the effect estimate: The true effect may be substantially different from the estimate of the effect

Very low certainty: Little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations

a. Clinically significant asthma exacerbations: episodes of asthma worsening with systemic corticosteroids for 3 or more days, a two-times increase in the dose of either inhaled corticosteroids or the need for asthma-related emergency treatment.

b. All included studies were funded by industry, and all showed positive results. No industry-independent observational or randomized studies were identified to contrast the results. Therefore, the quality of the evidence was downgraded for potential publication bias (70).

c. The pooled effect of risk ratio was assessed at 15 weeks (52) and the rate ratio was evaluated at 52 weeks (53)

d. Downgraded because the absolute effect includes both potential clinically meaningful benefits and harms.

e. Downgraded because FEV1 is considered a surrogate outcome of asthma control of symptoms, with a variable correlation with asthma symptoms (72).

f. The panel agreed that minimal important difference (MID) for FEV1 is 0.20 L g. Castro 2015 also assessed FEV1 at 52 weeks, the mean difference from baseline was 122.28 mL(45.54, 199.02). We also included 3 studies (52,54, 55) reporting FVC (mL), the pooled mean difference was 205.94 (88.69, 323.19); see full text report.

h. MID for ACQ-7 is 0.5 points (Juniper 2005).

i. Castro 2015 also assessed ACQ-7 at 52 weeks, the mean difference from baseline of ACQ-7 was -0.25 (-0.34, -0.16), see full text report.

j. MID for the Asthma symptoms utility index is an increase of 0.09 points (79).

k. MID of AQLQ is 0.5 points (37).

l. Castro 2015 also assessed AQLQ at 52 weeks, the mean difference from baseline was 0.29 (0.18, 0.41), see full text report.

m. Reduction of blood eosinophil counts is a surrogate endpoint and not validated as a valuable outcome for monitoring asthma therapy (80).

n. MID for rescue medication use is a reduction by 0.81 puffs/day (35).

o. The effect may both be harmful or beneficial.

p. Data regarding this outcome was extracted from www.clinicaltrials.gov for ref.52 and ref.53

q. Very few events in both arms, thus it is not possible to estimate precisely the effect size between arms